

Interference induced metallic-like behavior of a two-dimensional hole gas in asymmetric GaAs/In_xGa_{1-x}As/GaAs quantum well

G. M. Minkov,¹ A. V. Germanenko,² O. E. Rut,² A. A. Sherstobitov,¹ and B. N. Zvonkov³

¹*Institute of Metal Physics RAS, 620219 Ekaterinburg, Russia*

²*Institute of Physics and Applied Mathematics, Ural State University, 620083 Ekaterinburg, Russia*

³*Physical-Technical Research Institute, University of Nizhni Novgorod, 603600 Nizhni Novgorod, Russia*

(Dated: February 6, 2008)

The temperature and magnetic field dependences of the conductivity of the heterostructures with asymmetric In_xGa_{1-x}As quantum well are studied. It is shown that the metallic-like temperature dependence of the conductivity observed in the structures investigated is quantitatively understandable within the whole temperature range, $T = 0.4 - 20$ K. It is caused by the interference quantum correction at fast spin relaxation for $0.4 \text{ K} < T < 1.5$ K. At higher temperatures, $1.5 \text{ K} < T < 4$ K, it is due to the interaction quantum correction. Finally, at $T > 4 - 6$ K, the metallic-like behavior is determined by the phonon scattering.

PACS numbers: 73.20.Fz, 73.61.Ey

Transport properties of two dimensional (2D) systems reveal the intriguing features. One of them is a metallic-like temperature dependence of the conductivity, σ , at low temperature $d\sigma/dT < 0$. As a rule, such a behavior is observed in the structures with strong hole-hole (h - h) or electron-electron (e - e) interaction characterized by large value of the gas parameter $r_s = \sqrt{2}/(a_B k_F) \gg 1$, where a_B and k_F are the effective Bohr radius and the Fermi quasimomentum, respectively. That is why the e - e (h - h) interaction is considered as the main reason for the metallic-like T -dependence of σ .¹ The theory of the interaction quantum correction^{2,3} developed beyond the diffusive regime is widely applied to treat the experimental results.^{4,5}

On the other hand, there is a mechanism, which could in principle result in the metallic-like behavior of the system of weakly interacting electrons (holes).^{6,7,8,9} It is the interference quantum correction to the conductivity:¹⁰

$$\delta\sigma^{\text{WaL}} = G_0 \left\{ -\frac{1}{2} \ln \left(\frac{\tau}{\tau_\phi} \right) + \ln \left(\tau \left[\frac{1}{\tau_\phi} + \frac{1}{\tau_s} \right] \right) + \frac{1}{2} \ln \left(\tau \left[\frac{1}{\tau_\phi} + \frac{2}{\tau_s} \right] \right) \right\}, \quad G_0 = \frac{e^2}{2\pi^2 \hbar}, \quad (1)$$

where τ_s , τ_ϕ and τ are the spin, phase, and transport relaxation time, respectively. The sign of $d\sigma/dT$ for this mechanism depends on the relation between τ_s and τ_ϕ . When τ_s is appreciably shorter than τ_ϕ the interference correction is

$$\delta\sigma^{\text{WaL}} \simeq \frac{1}{2} G_0 \ln \frac{\tau_\phi}{\tau}. \quad (2)$$

Since the phase relaxation time at low temperatures is determined by electron-electron collisions, it is inversely proportional to the temperature, and, thus, the sign of $d\sigma/dT$ given by Eq. (2) is negative. In contrast to the weak localization characterized by the positive sign of $d\sigma/dT$, this phenomenon is known as the weak antilocalization (WaL). If the interference correction dominates the interaction correction, such a 2D system should demonstrate the metallic-like behavior.




FIG. 1: (Color online) The cross section and calculated valence band profile for the structure investigated at $V_g = 0$ V.

In this paper, we present results of the experimental study of the transport in the hole single quantum well with the asymmetric doping. Such a manner of doping results in the relatively large value of the spin-orbit splitting of the energy spectrum due to the Bychkov-Rashba mechanism¹¹ and, as consequence, in the relatively fast spin relaxation. We show experimentally that such a system really demonstrates the metallic-like behavior caused by the spin relaxation.

The heterostructure studied was grown by metal-organic vapor-phase epitaxy on a semi-insulator GaAs substrate. It consists of a 250 nm-thick undoped GaAs buffer layer, a 7 nm In_{0.2}Ga_{0.8}As well, a 7 nm spacer of undoped GaAs, a carbon δ -layer and 200 nm cap layer of undoped GaAs. Two samples from the same wafer have been studied. The samples were mesa etched into standard Hall bars. In order to tune the hole density, an Al gate electrode was deposited onto the cap layer. The hole density and mobility are $p = (4.0 - 7.5) \times 10^{11} \text{ cm}^{-2}$ and $\mu = (4000 - 8000) \text{ cm}^2/\text{V s}$ respectively. The va-




FIG. 2: The temperature dependence of the conductivity measured at different gate voltages. Symbols are the data, lines are the calculation results.

lence band profile calculated self-consistently is shown in Fig. 1. Similar results were obtained for both samples. Here we present the data obtained for one of them which parameters for some gate voltages are presented in Table I.

The temperature dependences of the conductivity measured for different gate voltages in the absence of a magnetic field are presented in Fig. 2. It is seen that they are metallic-like at the high hole density and mostly insulating at lower one. In order to understand the origin of such a behavior, the magnetic field dependences of the resistivity and the Hall effect has been investigated. The magnetoconductivity $\Delta\sigma(B) = \rho_{xx}^{-1}(B) - \rho_{xx}^{-1}(0)$ measured for one of the gate voltage, $V_g = 1.3\text{ V}$, at different temperatures are shown in Fig. 3(a). A characteristic antilocalization minimum is clearly seen; the lower is the temperature, the deeper is the minimum. As shown in Ref. 12, the Dyakonov-Perel mechanism¹³ is the main mechanism of spin relaxation in such a type systems. It has been also shown that the Bychkov-Rashba mechanism is responsible for the spin-orbit splitting of the energy spectrum in p -type GaAs/In_xGa_{1-x}As/GaAs quantum wells, and this splitting is cubic in quasimomentum. In this case the spin relaxation rate can be obtained from the fit of the magnetoconductivity curve by the Hikami-Larkin-Nagaoka expression.¹⁰ The results of the best fit

TABLE I: The parameters of structure for different gate voltages

V_g (V)	p (cm ⁻²)	τ (ps)	τ_ϕ (ps)	τ_s (ps)	F_0^σ	$\tilde{F}_0^\sigma$
0	6.5×10^{11}	0.73	$28/T$	4.8	-0.39	-0.34
1.3	4.8×10^{11}	0.48	$20/T$	11.0	-0.405	-0.36
1.6	4.4×10^{11}	0.39	$27/T$	17	-0.378	-0.322




FIG. 3: (a) – The $\Delta\sigma$ -vs- B dependence for $V_g = 1.3\text{ V}$. Solid lines are experimental data. Dashed lines are the results of the fitting procedure performed at $B < 0.2 B_{tr}$ (dashed lines) and $B < 0.1 B_{tr}$ (dotted lines). (b) and (c) – The T -dependence of τ_ϕ and τ_s obtained from the fit of the magnetoconductivity curves for $V_g = 0\text{ V}$ and 1.3 V , respectively at $B < 0.2 B_{tr}$ (solid symbols) and $B < 0.1 B_{tr}$ (open symbols), $B_{tr} = 91\text{ mT}$. Lines are the interpolating dependences which are used in calculation of $\delta\sigma^{\text{WaL}}$ and $\sigma(T)$ presented in Figs. 2 and 4.

are shown in Fig. 3(a). Since the theory¹⁰ was developed within the diffusion approximation, the fit was performed for the relatively low magnetic fields, $B < 0.1 B_{tr}$ and $B < 0.2 B_{tr}$, where $B_{tr} = \hbar/(2el^2)$ is the transport magnetic field, l is the transport mean free path.

The parameters τ_s and τ_ϕ found from the fit at different temperatures for $V_g = 0\text{ V}$ and 1.3 V are shown in Figs. 3(b) and 3(c), respectively. For $V_g = 0\text{ V}$, the data obtained from the different fitting range of magnetic field practically coincide. For $V_g = 1.3\text{ V}$, they are somewhat different that indicates an accuracy in determination of the phase and spin relaxation times. One can see that both τ_ϕ and τ_s demonstrate reasonable behavior. The τ_ϕ vs T dependence is close to T^{-1} -law that is typical for the 2D systems. Within the limits of experimental error the spin relaxation time τ_s is mostly independent of the temperature as should be in the degenerated gas of carriers.

The value of τ_s is significantly less than that of τ_ϕ only at $T \lesssim (1 - 3)\text{ K}$ depending on the gate voltage, that should provide the metallic-like temperature dependence of the conductivity only at the lowest temperatures. This is justified by the calculation results presented in Fig. 4. The dashed lines in this figure show the temperature dependence of the conductivity calculated as $\sigma(T) = \sigma_0 + \delta\sigma^{\text{WaL}}(T)$, where $\sigma_0 = e^2\tau n/m$ is the Drude conductivity determined in the given case by the ionized impurity scattering.¹⁴ The dependences $\delta\sigma^{\text{WaL}}(T)$ have been calculated from Eq. (1) with τ_s and $\tau_\phi(T)$ presented in Table I that well interpolate the corresponding experimental data shown in Figs. 3(b) and 3(c).

Thus, the weak antilocalization explain the negative sign of $d\sigma/dT$ only at lowest temperatures.

A different mechanism which can contribute to the temperature dependence of the conductivity is the h - h interaction. It can be conventionally subdivided into the ballistic and diffusion parts $\delta\sigma_b^{hh}$ and $\delta\sigma_d^{hh}$, respectively:^{2,15}

$$\delta\sigma^{hh}(T) = \delta\sigma_b^{hh}(T) + \delta\sigma_d^{hh}(T), \quad (3a)$$

$$\delta\sigma_b^{hh}(T) = 2\pi G_0 \frac{T\tau}{\hbar} \left[1 - \frac{3}{8} f(T\tau) \right] \quad (3b)$$

$$+ \frac{3\tilde{F}_0^\sigma}{1 + \tilde{F}_0^\sigma} \left(1 - \frac{3}{8} t(T\tau, \tilde{F}_0^\sigma) \right) \Bigg],$$

$$\delta\sigma_d^{hh}(T) = -G_0 \left[1 + 3 \left(1 - \frac{\ln(1 + F_0^\sigma)}{F_0^\sigma} \right) \right] \ln \frac{\hbar}{T\tau} \quad (3c)$$

where the functions $f(T\tau)$ and $t(T\tau, \tilde{F}_0^\sigma)$ are given in Ref. 2. The values of the Fermi-liquid constants F_0^σ and $\tilde{F}_0^\sigma$ (see Table I) have been found as described in Ref. 15, where the interaction correction was thoroughly studied in analogous systems.

Dotted and dash-dotted curves in Fig. 4 are drawn with taking into account only the diffusion and ballistics parts of the interaction correction, respectively. It is seen that their contributions to the temperature dependence are of different sign: $\delta\sigma_b^{hh}$ reveals the metallic like behavior whereas $\delta\sigma_d^{hh}$ is insulating. Thin solid lines in Fig. 4 are the total conductivity calculated as $\sigma(T) = \sigma_0 + \delta\sigma^{WaL}(T) + \delta\sigma^{hh}(T)$. It is seen that the consideration of both quantum corrections allows us to describe the experimental data in wider temperature range quantitatively. A good agreement is already evident up to $T \simeq 4 - 6$ K.

Finally, the decrease of the conductivity at higher temperature, $T \gtrsim 4 - 6$ K, is simply due to the phonon scattering, which cannot be ignored in heterostructures based on the piezoelectric crystals as our analysis shows. The dashed-double-dotted lines in Fig. 4 are the theoretical T -dependence of the Drude conductivity calculated taking into account the phonon scattering $\sigma_0^{ph} = \frac{e^2 n}{m} (\tau^{-1} + \tau_{ph}^{-1})^{-1}$, where τ_{ph}^{-1} is the phonon contribution to the scattering rate. The latter is calculated in Ref. 16 and represented in the form $\tau_{ph}(T) = a(T_0/T)^3 [1 + b(T/T_0)^2]$, where $T_0 = \sqrt{2mE_F S_t^2}$ with E_F and S_t as the Fermi energy and transverse sound velocity, respectively. The parameters a and b depend on the carriers parameters and crystal properties, and have been calculated according to Ref. 16 with $m = 0.16 m_0$.¹⁵ One can see that $\sigma_0^{ph}(T)$ really exhibits a considerably strong decrease explaining, thus, the experimental dependence $\sigma(T)$ at $T \gtrsim 4 - 6$ K.

The bold lines in Fig. 4 are calculated taking into account all three effects considered above: the interference quantum correction, the hole-hole interaction quantum correction, and the temperature dependence of the Drude




FIG. 4: (Color online) The T -dependence of σ for $V_g = 0$ V (a) and 1.3 V (b). Symbols are the experimental results. Lines are the results of calculation.

conductivity due to phonon scattering:

$$\sigma(T) = \sigma_0^{ph}(T) + \delta\sigma^{WaL}(T) + \delta\sigma^{hh}(T). \quad (4)$$

It is clearly seen that the calculation results are in excellent agreement with the experimental ones (see also Fig. 2).

Let us briefly discuss what type of the temperature dependence of the conductivity is expected at millikelvin temperatures. It is governed by competition between the interference and interaction quantum corrections. The interference correction $\delta\sigma^{WaL}$ at low temperatures has the form given by Eq. (2) and provides the logarithmic metallic-like behavior of $\sigma(T)$.

The diffusion part of the interaction correction $\delta\sigma_d^{hh}$ is logarithmic as well as $\delta\sigma^{WaL}$, but cannot be, however, described by Eq. (3c). The fact is that the second term in square brackets in Eq. (3c) is suppressed at low temperature, $T \ll \hbar/\tau_s$, so that the expression for $\delta\sigma_d^{hh}$ consists of only the Fock term^{8,18}

$$\delta\sigma^{hh}(T) \simeq \delta\sigma_d^{hh}(T) \simeq -G_0 \ln \frac{\hbar}{T\tau}, \quad T \ll \hbar/\tau_s. \quad (5)$$

In contrast to $\delta\sigma_d^{hh}$, the ballistic contribution of the interaction correction $\delta\sigma_b^{hh}$ is insensitive to the Bychkov-Rashba spin-relaxation mechanism,¹⁷ and vanishes at $T \rightarrow 0$ in accordance with Eq. (3). This is because the relevant ballistic paths contain two segments with opposite momenta, the spin rotation on the first segment is exactly canceled by counter-rotation of spin on the second segment.

Thus the total quantum correction $\delta\sigma^{WaL} + \delta\sigma^{hh}$ should be negative at very low temperatures, and the T -dependence of the conductivity is anticipated to be insulating-like

$$\sigma(T) \propto \delta\sigma^{WaL}(T) + \delta\sigma^{hh}(T) = -\frac{1}{2} G_0 \ln \frac{\hbar}{T\tau}. \quad (6)$$

Using the values of τ_s from Table I we estimate that this regime should happen in our samples at $T \ll 0.4 - 1.5$ K.

The experiments intended to verify this prediction are in our plan for the future.

In conclusion, we have experimentally studied the temperature dependence of the conductivity of 2D hole gas in asymmetrical GaAs/In_xGa_{1-x}As/GaAs quantum well heterostructures. It is shown that there is not one universal mechanism which is responsible for the metallic-like temperature dependence of the conductivity in the whole temperature range for the structures investigated. There are three physically different mechanisms: the weak antilocalization, the hole-hole interaction, and the phonon

scattering. Each of them is the main within the corresponding temperature range. The first one dominates at $T \simeq 0.4 - 1.5$ K. At higher temperatures, $T \simeq 1.5 - 4$ K, the interaction quantum correction becomes essential. Finally, the phonon scattering becomes important at $T \gtrsim 4 - 6$ K.

We are grateful to I. V. Gornyi for very useful discussions. This work was supported in part by the RFBR (Grant Nos. 05-02-16413, 06-02-16292, and 07-02-00528), the CRDF (Grant No. Y3-P-05-16), and by a Grant from the President of Russian Federation for Young Scientists.

-
- ¹ V. M. Pudalov, M. E. Gershenson, and H. Kojima, in *Fundamental Problems of Mesoscopic Physics: Interactions and Decoherence*, edited by I. V. Lerner, B. L. Altshuler, and Y. Gefen (Kluwer Academic Publishers, Dordrecht, 2004), p. 309; E. A. Galaktionov, A. K. Savchenko, S. S. Safonov, Y. Y. Proskuryakov, L. Li, M. Pepper, M. Y. Simmons, D. A. Ritchie, E. H. Linfield, and Z. D. Kvon, *ibid.* p. 349.
 - ² G. Zala, B. N. Narozhny, and I. L. Aleiner, *Phys. Rev. B* **64**, 214204 (2001).
 - ³ I. V. Gornyi and A. D. Mirlin, *Phys. Rev. Lett.* **90**, 076801 (2003); *Phys. Rev. B* **69**, 045313 (2004).
 - ⁴ Y. Y. Proskuryakov, A. K. Savchenko, S. S. Safonov, M. Pepper, M. Y. Simmons, and D. A. Ritchie, *Phys. Rev. Lett.* **89**, 076406 (2002).
 - ⁵ V. M. Pudalov, M. E. Gershenson, H. Kojima, G. Brunthaler, A. Prinz, and G. Bauer, *Phys. Rev. Lett.* **91**, 126403 (2003).
 - ⁶ M. A. Skvortsov, *Pis'ma ZhETF* **67**, 118 (1998) [*JETP Lett* **67**, 133 (1998)].
 - ⁷ Yuli Lyanda-Geller, *Phys. Rev. Lett.* **80**, 4273 (1998).
 - ⁸ I. V. Gornyi, A. P. Dmitriev, and V. Yu. Kachorovskii, *Pis'ma ZhETF* **68**, 314 (1998) [*JETP Lett* **68**, 338 (1998)].
 - ⁹ L. E. Golub and S. Pedersen, *Phys. Rev. B* **65**, 245311 (2002).
 - ¹⁰ S. Hikami, A. I. Larkin, and Y. Nagaoka, *Prog. Theor. Phys.* **63**, 707 (1980).
 - ¹¹ Y. A. Bychkov and E. I. Rashba, *J. Phys. C* **17**, 6039 (1984).
 - ¹² G. M. Minkov, A. A. Sherstobitov, A. V. Germanenko, O. E. Rut, V. A. Larionova, and B. N. Zvonkov, *Phys. Rev. B* **71**, 165312 (2005).
 - ¹³ M. I. Dyakonov and V. I. Perel, *Zh. Eksp. Teor. Fiz.* **60**, 1954 (1971) [*Sov. Phys. JETP* **33**, 1053 (1971)].
 - ¹⁴ More precisely, σ_0 includes also the temperature independent contributions of the interaction correction (see Ref. 2 for details), which are omitted in Eq. (3). The value of σ_0 has been experimentally obtained as described by G. M. Minkov, A. V. Germanenko, O. E. Rut, A. A. Sherstobitov, and B. N. Zvonkov in cond-mat/0606566. It should be stressed, however, that we do not need to know the value of σ_0 with the high accuracy to interpret the results presented here. In particular the calculation results presented in Fig. 2 are insensitive to σ_0 at all.
 - ¹⁵ G. M. Minkov, A. A. Sherstobitov, A. V. Germanenko, O. E. Rut, V. A. Larionova, B. N. Zvonkov, *Phys. Rev. B* **72**, 165325 (2005).
 - ¹⁶ V. Karpus, *Semicond. Sci. Technol.* **5**, 691 (1990).
 - ¹⁷ A. P. Dmitriev, I. V. Gornyi, V. Yu. Kachorovskii, and I. S. Lyubinskiy, unpublished; private communication.
 - ¹⁸ B. L. Altshuler and A. G. Aronov, *Solid State Comm.* **46**, 429 (1983).